



OPTIONS FOR UPDATING REGIMENS TO INCLUDE **KISUNLA**

INSTRUCTIONS FOR THE ORACLE HEALTH ELECTRONIC HEALTH RECORD (EHR) SYSTEM

Please refer to [page 4](#) for the updated Kisunla dosing schedule.

INDICATION AND USAGE

Kisunla is indicated for the treatment of Alzheimer's disease (AD). Treatment with Kisunla should be initiated in patients with mild cognitive impairment (MCI) or mild dementia stage of disease, the population in which treatment was initiated in the clinical trials.

SELECT IMPORTANT SAFETY INFORMATION

WARNING: AMYLOID-RELATED IMAGING ABNORMALITIES

Monoclonal antibodies directed against aggregated forms of beta amyloid, including Kisunla, can cause amyloid-related imaging abnormalities (ARIA), characterized as ARIA with edema (ARIA-E) and ARIA with hemosiderin deposition (ARIA-H). ARIA can be serious and life threatening events can occur. ARIA can be fatal. Serious intracerebral hemorrhages >1 cm, some fatal, have been observed with this class of medications. Because ARIA-E can cause focal neurologic deficits that can mimic an ischemic stroke, treating clinicians should consider whether such symptoms could be due to ARIA-E before giving thrombolytic therapy.

Apolipoprotein E $\epsilon 4$ (ApoE $\epsilon 4$) Homozygotes: Patients treated with this class of medications, including Kisunla, who are ApoE $\epsilon 4$ homozygotes have a higher incidence of ARIA, including symptomatic and serious ARIA, compared to heterozygotes and noncarriers. Testing for ApoE $\epsilon 4$ status should be performed prior to initiation of treatment to inform the risk of developing ARIA. Prior to testing, the risk of ARIA across genotypes and the implications of genetic testing results should be discussed with patients.

Consider the benefit for the treatment of Alzheimer's disease and risk of ARIA when deciding to treat with Kisunla.

Please see additional Important Safety Information on [pages 14-16](#), and full [Prescribing Information](#), including Boxed Warning regarding ARIA, and [Medication Guide](#) for Kisunla.

CONTENTS

1. Overview and Limitations	3
2. Indication and Usage	3
3. Updated Kisunla™ Dosing Schedule and Oracle Health EHR Implementation Guidance	4
4. Dosing and Administration	5
Patient Selection	5
Dosing Instructions	5
Monitoring and Dosing Interruption for Amyloid-Related Imaging Abnormalities	6
Dilution Instructions	8
Administration Instructions	9
5. Oracle Health Electronic Health Record Instructions	10
Updating the PowerPlan	10
Building the Regimen	12
6. Disclaimers	13
7. IMPORTANT SAFETY INFORMATION FOR Kisunla™ (donanemab-azbt)	14
8. Notes	17

Please see additional Important Safety Information on [pages 14-16](#), and full [Prescribing Information](#), including Boxed Warning regarding ARIA, and [Medication Guide](#) for Kisunla.

1. Overview and Limitations

Regimens contain orders needed to plan the patient's care for the duration of their therapy, from pre-infusion orders, during infusion assistance orders, to post-infusion needs, such as maintenance medications and imaging. PowerPlans define the cycles within the Regimen and enable the proper flow of the treatments. The Regimen Builder can be used to combine multiple PowerPlans into a single Regimen. Regimens enable date adjustments to persist for all PowerPlans in the Regimen.

After initial release, Regimens and PowerPlans may benefit from a clinical update. The optimization of these Regimens in the Oracle Health EHR system is a standard process and provides an opportunity to incorporate treatment updates. A customer may choose to update existing Regimens and their supporting PowerPlans to include additional treatments, such as Kisunla™ (donanemab-azbt).

This document is intended to provide health systems with instructions on how to update a Regimen with Kisunla, within the approved indication and consistent with the full Prescribing Information. These instructions are specific to the Oracle Health EHR system and to the use of Kisunla in the treatment of early symptomatic Alzheimer's disease and are not appropriate for other conditions, treatments, therapeutic areas, or for other EHR systems. These instructions specifically detail how to add dosing information and infusion instructions for Kisunla.

The process outlined on the following pages is variable, and not all steps will apply to every customer. Any steps or settings outlined in this document that are not part of a customer's standard process should be excluded or modified accordingly. Any questions should be directed to the appropriate service provider. The customer is solely responsible for implementing, testing, monitoring, and maintaining the ongoing operation of any EHR resources. The elements provided herein are only suggestions, and it is strongly recommended that clinical and operational leadership determines that the final elements align with the expectations and goals of the organization. A Notes section (located on [page 17](#) of this guide) can be used to document any additions or changes that need to be made prior to implementation.

Typically, a health system will conduct a clinical review process to confirm and approve the suggested build and subsequent optimizations. Various stakeholders may participate in reviewing Regimen and PowerPlan requests prior to implementation.

This document is not intended to provide any clinical advice or recommendations, which are solely the responsibility of the customer and its healthcare providers. Treatment selection is always a decision made by the healthcare provider, and Regimens may be overridden to reflect this. An EHR newsletter or other communication medium may be considered to notify end users of the availability and contents of any updated Regimens.

2. Indication and Usage

Kisunla is indicated for the treatment of Alzheimer's disease (AD). Treatment with Kisunla should be initiated in patients with mild cognitive impairment (MCI) or mild dementia stage of disease, the population in which treatment was initiated in the clinical trials.

SELECT IMPORTANT SAFETY INFORMATION

- **Hypersensitivity: Kisunla is contraindicated** in patients with known serious hypersensitivity to donanemab-azbt or to any of the excipients. Reactions have included anaphylaxis.

Please see additional Important Safety Information on [pages 14-16](#), and full [Prescribing Information](#), including Boxed Warning regarding ARIA, and [Medication Guide](#) for Kisunla.



3. Updated Kisunla™ Dosing Schedule and Oracle Health EHR Implementation Guidance

The dosing schedule for Kisunla (donanemab-azbt) has been updated, impacting how it is prescribed to patients. The new dosing schedule is as follows:

- **Infusion 1:** 350 mg administered over approximately 30 minutes.
- **Infusion 2:** 700 mg administered over approximately 30 minutes.
- **Infusion 3:** 1050 mg administered over approximately 30 minutes.
- **Infusion 4 and beyond:** 1400 mg administered over approximately 30 minutes

Please refer to the information within the Dosing and Administration section of this guide for additional details. Organizations must determine their course of action based on whether they have already implemented PowerPlans and Regimens for Kisunla in their Oracle Health EHR system:

For Organizations With an Existing Regimen for Kisunla:

- **Update Existing Regimen for Kisunla:** Based on your organizational protocols, review and update the current PowerPlans and Regimen build for Kisunla to align with the new dosing schedule. This includes modifying and implementing additional PowerPlans to ensure all doses are accounted for and adding them to the existing Regimen build.

For Organizations Without an Existing Regimen for Kisunla:

- **Follow the Instructions to Update an Existing Regimen for Kisunla:** Refer to your organization's standard process for implementing PowerPlans and Regimens within your Oracle Health EHR system, ensuring all stakeholders are informed and trained on the new protocol. Refer to [section 5](#) of this guide for Oracle Health EHR system instructions.

SELECT IMPORTANT SAFETY INFORMATION

Amyloid-Related Imaging Abnormalities (ARIA)

ARIA usually occurs early in treatment and is usually asymptomatic, although serious and life-threatening events, including seizure and status epilepticus, can occur. ARIA can be fatal. When present, reported symptoms associated with ARIA may include, but are not limited to, headache, confusion, visual changes, dizziness, nausea, and gait difficulty. Focal neurologic deficits may also occur. Symptoms associated with ARIA usually resolve over time.

Please see additional Important Safety Information on [pages 14-16](#), and full [Prescribing Information](#), including Boxed Warning regarding ARIA, and [Medication Guide](#) for Kisunla.



4. Dosing and Administration

The following is an overview and quick-reference guide for the dosing and administration details for Kisunla™ for its approved indication as per the full Prescribing Information. Step-by-step instructions for applying some of these medication details in the Regimen update are shown in the following section of this guide.

Patient Selection

Confirm the presence of amyloid beta pathology prior to initiating treatment [see Clinical Pharmacology (12.1)]. Obtain a recent baseline brain magnetic resonance imaging (MRI) prior to initiating treatment with Kisunla.

Dosing Instructions

The recommended dosage of Kisunla is 350 mg for the first dose, 700 mg for the second dose, 1050 mg for the third dose (350/700/1050 mg), followed by 1400 mg every four weeks (see Table 1 in the full Prescribing Information, reproduced below). Kisunla is administered every four weeks as an intravenous infusion over approximately 30 minutes. Kisunla must be diluted prior to administration (see Table 4 in the full Prescribing Information, reproduced on [page 8](#)).

Table 1 From Full Prescribing Information: Dosing Schedule

Intravenous Infusion (every 4 weeks)	Kisunla Dosage (administered over approximately 30 minutes)
Infusion 1	350 mg
Infusion 2	700 mg
Infusion 3	1050 mg
Infusion 4 and beyond	1400 mg

Consider stopping dosing with Kisunla based on reduction of amyloid plaques to minimal levels on amyloid PET imaging. In Study 1 and Study 2, dosing was stopped based on a reduction of amyloid levels below predefined thresholds on PET imaging [see *Clinical Studies* (14)].

If an infusion is missed, resume administration every four weeks at the same dose as soon as possible.

SELECT IMPORTANT SAFETY INFORMATION

Risk Factors for ARIA and Intracerebral Hemorrhages (ICH)

- **ApoE ε4 Carrier Status:** The risk of ARIA, including symptomatic and serious ARIA, is increased in apolipoprotein E ε4 (ApoE ε4) homozygotes.
- Recommendations for management of ARIA do not differ based on ApoE ε4 carrier status. Testing for ApoE ε4 status should be performed prior to initiation of treatment to inform the risk of developing ARIA.

→ See next page for additional content.

Please see additional Important Safety Information on [pages 14-16](#), and full [Prescribing Information](#), including Boxed Warning regarding ARIA, and [Medication Guide](#) for Kisunla.



4. Dosing and Administration (Continued)

Monitoring and Dosing Interruption for Amyloid-Related Imaging Abnormalities

Monitoring for ARIA

Obtain a recent baseline brain magnetic resonance imaging (MRI) prior to initiating treatment with Kisunla. Obtain an MRI prior to the 2nd, 3rd, 4th, and 7th infusions. If a patient experiences symptoms suggestive of ARIA, clinical evaluation should be performed, including an MRI if indicated.

Recommendations for Dosing Interruptions in Patients With ARIA

ARIA-E

The recommendations for dosing interruptions for patients with ARIA-E are provided in Table 2 of the full Prescribing Information, reproduced below:

Table 2 From Full Prescribing Information: Dosing Recommendations for Patients With ARIA-E¹

	ARIA-E Severity on MRI		
Clinical Symptom Severity ^a	Mild	Moderate	Severe
Asymptomatic	May continue dosing at current dose and schedule	Suspend dosing ^b	Suspend dosing ^b
Mild	May continue dosing based on clinical judgment	Suspend dosing ^b	
Moderate or Severe	Suspend dosing ^b		

^aMild: discomfort noticed, but no disruption of normal daily activity; Moderate: discomfort sufficient to reduce or affect normal daily activity; Severe: incapacitating, with inability to work or to perform normal daily activity.

^bSuspend until MRI demonstrates radiographic resolution and symptoms, if present, resolve; consider a follow-up MRI to assess for resolution 2 to 4 months after initial identification. Resumption of dosing should be guided by clinical judgment.

ARIA-E=amyloid-related imaging abnormalities-edema; MRI=magnetic resonance imaging.

SELECT IMPORTANT SAFETY INFORMATION

Risk Factors for ARIA and Intracerebral Hemorrhages (ICH) (Continued)

- **Radiographic Findings of Cerebral Amyloid Angiopathy (CAA):** Neuroimaging findings that may indicate CAA include evidence of prior ICH, cerebral microhemorrhage, and cortical superficial siderosis. CAA has an increased risk for ICH. The presence of an ApoE ε4 allele is also associated with CAA.

→ See next page for additional content.

Please see additional Important Safety Information on [pages 14-16](#), and full [Prescribing Information](#), including Boxed Warning regarding ARIA, and [Medication Guide](#) for Kisunla.



4. Dosing and Administration (Continued)

Monitoring and Dosing Interruption for Amyloid-Related Imaging Abnormalities (Continued)

Recommendations for Dosing Interruptions in Patients With ARIA (Continued)

ARIA-H

The recommendations for dosing interruptions for patients with ARIA-H are provided in Table 3 of the full Prescribing Information, reproduced below:

Table 3 From Full Prescribing Information: Dosing Recommendations for Patients With ARIA-H¹

	ARIA-H Severity on MRI		
Clinical Symptom Severity	Mild	Moderate	Severe
Asymptomatic	May continue dosing at current dose and schedule	Suspend dosing ^a	Suspend dosing ^b
Symptomatic	Suspend dosing ^a		

^aSuspend until MRI demonstrates radiographic stabilization and symptoms, if present, resolve; resumption of dosing should be guided by clinical judgment; consider a follow-up MRI to assess for stabilization 2 to 4 months after initial identification.

^bSuspend until MRI demonstrates radiographic stabilization and symptoms, if present, resolve. Use clinical judgment when considering whether to continue treatment or permanently discontinue Kisunla.

ARIA-H=amyloid-related imaging abnormalities-hemosiderin deposition.

In patients who develop intracerebral hemorrhage greater than 1 cm in diameter during treatment with Kisunla, suspend dosing until MRI demonstrates radiographic stabilization and symptoms, if present, resolve. Resumption of dosing should be guided by clinical judgment.¹

SELECT IMPORTANT SAFETY INFORMATION

Risk Factors for ARIA and Intracerebral Hemorrhages (ICH) (Continued)

- The baseline presence of at least 2 microhemorrhages or the presence of at least 1 area of superficial siderosis on MRI, which may be suggestive of CAA, were identified as risk factors for ARIA. Patients were excluded from enrollment for findings on neuroimaging of prior ICH >1 cm in diameter, >4 microhemorrhages, >1 area of superficial siderosis, severe white matter disease, and vasogenic edema.

→ See next page for additional content.

Please see additional Important Safety Information on [pages 14-16](#), and full [Prescribing Information](#), including Boxed Warning regarding ARIA, and [Medication Guide](#) for Kisunla.



4. Dosing and Administration (Continued)

Dilution Instructions

- Prior to administration, Kisunla must be diluted with 0.9% sodium chloride injection, USP.
- Use aseptic technique when preparing the diluted Kisunla solution for intravenous infusion.
- Allow Kisunla to equilibrate to room temperature before preparation.
- Parenteral drug products should be inspected visually for particulate matter and discoloration prior to administration, whenever solution and container permit. Kisunla solution is clear to opalescent, colorless to slightly yellow to slightly brown. Do not use if particulate matter or discolorations are present.
- Withdraw required volume of Kisunla and mix with 0.9% sodium chloride injection, to the recommended total volume for a final concentration of 4 mg/mL to 10 mg/mL (see Table 4 in the full Prescribing Information, reproduced below). Use only 0.9% sodium chloride injection for dilution.

Table 4 From Full Prescribing Information: Preparation of Kisunla

Kisunla Dose (mg)	Kisunla Volume (mL)	Volume of 0.9% Sodium Chloride Injection Diluent (mL)	Final Volume of Diluted Solution to be Infused (mL)	Final Concentration of Diluted Solution (mg/mL) ^a
350 mg	20 mL	15 mL to 67.5 mL	35 mL to 87.5 mL	350 mg/87.5 mL (4 mg/mL) to 350 mg/35 mL (10 mg/mL)
700 mg	40 mL ^b	30 mL to 135 mL	70 mL to 175 mL	700 mg/175 mL (4 mg/mL) to 700 mg/70 mL (10 mg/mL)
1050 mg	60 mL ^c	45 mL to 202.5 mL	105 mL to 262.5 mL	1050 mg/262.5 mL (4 mg/mL) to 1050 mg/105 mL (10 mg/mL)
1400 mg	80 mL ^d	60 mL to 270 mL	140 mL to 350 mL	1400 mg/350 mL (4 mg/mL) to 1400 mg/140 mL (10 mg/mL)

^aFinal concentration of 4 mg/mL to 10 mg/mL.

^b2 vials of Kisunla.

^c3 vials of Kisunla.

^d4 vials of Kisunla.

SELECT IMPORTANT SAFETY INFORMATION

Risk Factors for ARIA and Intracerebral Hemorrhages (ICH) (Continued)

- **Concomitant Antithrombotic or Thrombolytic Medication:** In Study 1, the majority of exposures to antithrombotic medications were to aspirin. The number of events and the limited exposure to non-aspirin antithrombotic medications limit definitive conclusions about the risk of ARIA or ICH in patients taking antithrombotic medications. Additional caution should be exercised when considering the administration of antithrombotics or a thrombolytic agent (eg, tissue plasminogen activator) to a patient already being treated with Kisunla. One fatal ICH occurred in a patient taking Kisunla in the setting of focal neurologic symptoms of ARIA and the use of a thrombolytic agent in Study 1, and one fatal intracerebral hemorrhage occurred in the setting of ARIA and the use of a thrombolytic agent in Study 2.

Please see additional Important Safety Information on [pages 14-16](#), and full [Prescribing Information](#), including Boxed Warning regarding ARIA, and [Medication Guide](#) for Kisunla.



4. Dosing and Administration (Continued)

Dilution Instructions (Continued)

- Each vial is for one-time use only. Discard any unused portion left in the vial.
- Gently invert the diluted Kisunla solution to mix completely. Do not shake.
- After dilution, immediate use is recommended (see Description [Section 11] in the full Prescribing Information). If the diluted Kisunla solution is not administered immediately, store refrigerated at 2 °C to 8 °C (36 °F to 46 °F) for up to 72 hours or at room temperature (20 °C to 25 °C [68 °F to 77 °F]) for up to 12 hours.
- Do not freeze the diluted Kisunla solution.
- Storage times include the duration of infusion.

Administration Instructions

- Visually inspect the diluted Kisunla solution for particles or discoloration prior to administration. Do not use if it is discolored, or opaque, or foreign particles are seen.
- Prior to infusion, if the diluted solution has been stored under refrigeration, allow the diluted Kisunla solution to warm to room temperature.
- Administer the entire diluted solution intravenously over approximately 30 minutes.
- Promptly discontinue the infusion upon the first observation of any signs or symptoms consistent with a hypersensitivity-type reaction (see Warnings and Precautions [Section 5.2] in the full Prescribing Information).
- Flush the line only with 0.9% sodium chloride injection at the end of the infusion per access-specific line maintenance protocol.
- Observe the patient post-infusion for a minimum of 30 minutes to evaluate for infusion reactions and hypersensitivity reactions (see Warnings and Precautions [Section 5.2] in the full Prescribing Information).

SELECT IMPORTANT SAFETY INFORMATION

Risk Factors for ARIA and Intracerebral Hemorrhages (ICH) (Continued)

- Consider whether ischemic stroke symptoms could be due to ARIA-E before giving thrombolytic therapy in a patient being treated with Kisunla, because ARIA-E can cause focal neurologic deficits that can mimic an ischemic stroke.
- Caution should be exercised when considering the use of Kisunla in patients with factors that indicate an increased risk for ICH and in particular for patients who need to be on anticoagulant therapy or patients with findings on MRI that are suggestive of CAA.

Please see additional Important Safety Information on [pages 14-16](#), and full [Prescribing Information](#), including Boxed Warning regarding ARIA, and [Medication Guide](#) for Kisunla.



5. Oracle Health Electronic Health Record Instructions

Confirm a diagnosis of early symptomatic Alzheimer's disease (mild cognitive impairment or mild dementia stage of disease) and also the presence of amyloid beta pathology prior to initiating treatment (See Patient Selection [section 2.1] In the full Prescribing Information).

According to the organization's guidelines, build PowerPlans based on treatment cycles. Use the Regimen Builder to group the PowerPlans into a single-treatment Regimen.

- Select an existing PowerPlan that can be updated to include Kisunla or if there is no appropriate PowerPlan for Kisunla, consider creating a new PowerPlan*.
- Update either a single PowerPlan or multiple PowerPlans using the PowerPlan tool.
- Update “oncology-like” plans based on medication(s) and treatment cycle.
- Include orders for medication administration, associated labs, imaging, administration instructions, etc.
- Use the Cycle feature of each PowerPlan to define required cycles of treatment.
- Use the Regimen Builder to include all appropriate PowerPlans in a single-treatment Regimen.

5.1 Updating the PowerPlan

This guide assumes four PowerPlan builds, one for each Kisunla dose (350 mg, 700 mg, 1050 mg, and 1400 mg). Each PowerPlan will be included in the Regimen builder according to the number of cycles each requires. Cycle 1 will be for the 350 mg dose, cycle 2 will be for the 700 mg dose, cycle 3 will be for the 1050 mg dose, and cycle 4 and beyond will be for the 1400 mg dose.

Note: A full year of treatment may require either 12 or 13 total cycles; this guide assumes a 12 treatment per year scenario.

1. From the Millennium applications list, select **DCPTools.exe.**; under **Order Management**, select **PowerPlan Tool**.
2. Update to a new PowerPlan and name based on organizations naming convention; for example:
 - a. **Name:** Kisunla 350 mg.
 - b. **Phase:** Multiple.

SELECT IMPORTANT SAFETY INFORMATION

ARIA Monitoring and Dose Management Guidelines

- Baseline brain MRI and periodic monitoring with MRI are recommended prior to the 2nd, 3rd, 4th, and 7th infusions. Enhanced clinical vigilance for ARIA is recommended during the first 24 weeks of treatment with Kisunla. If a patient experiences symptoms suggestive of ARIA, clinical evaluation should be performed, including MRI if indicated. If ARIA is observed on MRI, careful clinical evaluation should be performed prior to continuing treatment.

Please see additional Important Safety Information on [pages 14-16](#), and full [Prescribing Information](#), including Boxed Warning regarding ARIA, and [Medication Guide](#) for Kisunla.



5. Oracle Health Electronic Health Record Instructions (Continued)

Updating the PowerPlan (Continued)

3. Enter details:
 - a. **Type:** For example, Medical (according to the organization protocol).
 - b. **Display Method:** For example, select Clinical Category (to display phases by category, such as Medications, Imaging, etc.) or use descriptive names, such as Pretreatment Orders.
 - c. **Status:** Testing.
 - d. **Name the Phases:** For example, Pretreatment Orders, Imaging Orders, Medication Administration Orders.
4. From the **Attributes** section on the right side of the window, **Cycle Settings** option, select **Click here** to review.
 - a. Define the Cycles as Range, enter the desired number of cycles; for example, 1 for Kisunla 350 mg.

Note: Each time this PowerPlan is ordered, it will automatically default to the cycle next in line to be used. When building the 1400 mg PowerPlan the Cycle Range will be 4-12.
5. To add an order, click on the phase, such as **Pretreatment Orders** phase; then, from the lower right pane, the **Orders** tab, add an order for **Brain MRI W W/O Contrast**.

Note: the MRI will also be included in the Pretreatment Orders phase of cycles 2, 3, 4, and 7.
6. Highlight the **Medication Administration Orders** phase. Then from the **Orders** tab, search for and select **Kisunla 350 mg**; select **Add**.
7. Highlight the **Kisunla 350 mg** and from the right-side pane, select the **Order Sentence** tab.

*If creating a new Order Group, the system should consider what other medications may also warrant inclusion.

SELECT IMPORTANT SAFETY INFORMATION

ARIA Monitoring and Dose Management Guidelines (Continued)

Recommendations for dosing in patients with ARIA-E and ARIA-H depend on clinical symptoms and radiographic severity. Depending on ARIA severity, use clinical judgment in considering whether to continue dosing, interrupt treatment, or permanently discontinue Kisunla. See Prescribing Information for additional dosing considerations.

Hypersensitivity Reactions

Hypersensitivity reactions, including anaphylaxis and angioedema, have occurred in patients who were treated with Kisunla. Promptly discontinue the infusion upon the first observation of any signs or symptoms consistent with a hypersensitivity reaction and initiate appropriate therapy.

Please see additional Important Safety Information on [pages 14-16](#), and full [Prescribing Information](#), including Boxed Warning regarding ARIA, and [Medication Guide](#) for Kisunla.



5. Oracle Health Electronic Health Record Instructions (Continued)

Updating the PowerPlan (Continued)

8. From the list of prebuilt order sentences, select **Properly dilute and administer 350 mg over 30 minutes. Observe the patient post-infusion for a minimum of 30 minutes to evaluate for infusion reactions and hypersensitivity reactions.**
[See Section 5.3, Infusion-Related Reactions, in the full Prescribing Information].
*Note: If prebuilt order sentences have not been built, click on the **Order Sentence** box at the bottom of the **Order Details** section to add.*
9. Input order details, such as **Dose, Dose Unit, Route of Administration, Frequency**, etc.
10. After all desired orders have been built in a phase, highlight the phase name, from the **Attributes** section, **Treatment Schedule** option, select **Click here** to review.
11. In the **Build Treatment Schedule – Treatment** window, set up the schedule for the medications.
 - a. Enter the treatment **Schedule Length**; for example, 28 days and the **Start With** number, such as 1.
 - b. Set the **Day of Treatment** values, such as duration and unit; for example, 12 hours for outpatient orders.
 - c. Link all orders to the treatment schedule; for example, Kisunla 350 mg link to Day 1.

Duplicate these steps to update a PowerPlan for remaining Kisunla doses (700 mg, 1050 mg, and 1400 mg).

5.2 Building the Regimen

1. From the **Regimen Builder**, select **New Regimen**; enter the Regimen name, such as Kisunla Regimen, and a description.
2. Leave the new Regimen active and fill in details according to organization protocols.
3. Select **+Add**; add the previously updated PowerPlans; add the 350 mg, 700 mg, and 1050 mg PowerPlan once each; add the 1400 mg PowerPlan 9 times.
4. Highlight each PowerPlan, choose **Cycle Settings**; select the appropriate **Default Plan Cycle** number for each.
5. Then highlight the Regimen (name), from the **Detail** options, **Offsets**, select **Click here** to add **Regimen Offsets**.
6. Enter the amount of Offset and the **Offset Unit** needed between cycles; for example, 4 weeks.
7. Enter the **Anchor Element between Cycles**; for example, Cycle 1 is the Anchor Element for Cycle 2 and so on.
8. Click **OK** to close the Regimen Offsets, then click **Save** to save the Regimen.

SELECT IMPORTANT SAFETY INFORMATION

Infusion-Related Reactions (IRR)

IRRs were observed with Kisunla with the majority occurring within the first 4 infusions. Most IRRs occurred during the infusion or within 30 minutes after completion of the infusion, however some have occurred hours after an infusion. Signs and symptoms of IRRs include chills, erythema, nausea/vomiting, flushing, difficulty breathing/dyspnea, sweating, elevated blood pressure, headache, chest pain, and low blood pressure. In the event of an IRR, the infusion rate may be reduced, or the infusion may be discontinued, and appropriate therapy initiated as clinically indicated. Consider pre-treatment with antihistamines, acetaminophen, or corticosteroids prior to subsequent dosing.

Please see additional Important Safety Information on [pages 14-16](#), and full [Prescribing Information](#), including Boxed Warning regarding ARIA, and [Medication Guide](#) for Kisunla.



6. Disclaimers

- The Customer (eg, the physician, medical group, integrated delivery network [IDN]) shall be solely responsible for the implementation, testing, and monitoring of the instructions to ensure proper orientation in each Customer's EHR system.
- Capabilities, functionality, and set-up (customization) for each EHR system may vary. Lilly shall not be responsible for revising the implementation instructions it provides to any Customer if the Customer modifies or changes its software, or the configuration of its EHR system, after such time as the implementation instructions have been initially provided by Lilly.
- While Lilly tests its implementation instructions on multiple EHR systems, the instructions are not guaranteed to work for all available EHR systems, and Lilly shall have no liability thereto.
- While EHRs may assist providers in identifying appropriate patients for consideration of assessment, treatment, and referral, the decision and action should ultimately be decided by a provider in consultation with the patient, after a review of the patient's records to determine eligibility, and Lilly shall have no liability related to a provider's decision and action (or inaction) regarding any patient identified or treated using this resource.
- The instructions have not been designed to and are not resources and/or solutions for meeting Advancing Care Information and/or any other quality/accreditation requirement.
- All product/company names mentioned herein are the trademarks of their respective owners, all rights reserved. Reference to these products is not intended to imply affiliation with or sponsorship of Lilly and/or its affiliates.

Please see additional Important Safety Information on [pages 14-16](#), and full [Prescribing Information](#), including Boxed Warning regarding ARIA, and [Medication Guide](#) for Kisunla.



7. IMPORTANT SAFETY INFORMATION FOR Kisunla™ (donanemab-azbt)

WARNING: AMYLOID-RELATED IMAGING ABNORMALITIES

Monoclonal antibodies directed against aggregated forms of beta amyloid, including Kisunla, can cause amyloid-related imaging abnormalities (ARIA), characterized as ARIA with edema (ARIA-E) and ARIA with hemosiderin deposition (ARIA-H). ARIA usually occurs early in treatment and is usually asymptomatic, although serious and life-threatening events can occur. ARIA can be fatal. Serious intracerebral hemorrhages >1 cm, some of which have been fatal, have been observed in patients treated with this class of medications. Because ARIA-E can cause focal neurologic deficits that can mimic an ischemic stroke, treating clinicians should consider whether such symptoms could be due to ARIA-E before giving thrombolytic therapy in a patient being treated with Kisunla.

ApoE ε4 Homozygotes: Patients treated with this class of medications, including Kisunla, who are apolipoprotein E ε4 (ApoE ε4) homozygotes (approximately 15% of Alzheimer's disease patients) have a higher incidence of ARIA, including symptomatic, serious, and severe radiographic ARIA, compared to heterozygotes and noncarriers. Testing for ApoE ε4 status should be performed prior to initiation of treatment to inform the risk of developing ARIA. Prior to testing, the risk of ARIA across genotypes and the implications of genetic testing results should be discussed with patients.

Consider the benefit for treating Alzheimer's disease and risk of ARIA when deciding to treat with Kisunla.

Kisunla is **contraindicated** in patients with known serious hypersensitivity to donanemab-azbt or to any of the excipients. Reactions have included anaphylaxis.

Amyloid-Related Imaging Abnormalities (ARIA)

ARIA usually occurs early in treatment and is usually asymptomatic, although serious and life-threatening events, including seizure and status epilepticus, can occur. ARIA can be fatal. When present, reported symptoms associated with ARIA may include, but are not limited to, headache, confusion, visual changes, dizziness, nausea, and gait difficulty. Focal neurologic deficits may also occur. Symptoms associated with ARIA usually resolve over time.

In Study 1, safety was assessed in patients who received Kisunla Dosing Regimen 1 (n=853) compared to those who received placebo (n=874). In Study 2, the effect of different dosing regimens of Kisunla on ARIA was assessed, including in patients who received Kisunla Dosing Regimen 2 (n=212).

Incidence of ARIA

A lower incidence of ARIA was observed with Dosing Regimen 2 as compared to Dosing Regimen 1. Therefore, Dosing Regimen 2 is the recommended dosage for Kisunla.

In Study 1, symptomatic ARIA-E occurred in 6% of patients through 18 months of treatment with Kisunla.

Clinical symptoms associated with ARIA resolved in approximately 85% of those patients.

Including asymptomatic radiographic events, ARIA, ARIA-E, and ARIA-H were observed with Kisunla: 36%, 24%, and 31% of patients treated with Kisunla, respectively compared to 14%, 2%, and 13% of patients on placebo. There was no increase in isolated ARIA-H for Kisunla vs placebo.

Please see additional Important Safety Information on [pages 15-16](#), and full [Prescribing Information](#), including Boxed Warning regarding ARIA, and [Medication Guide](#) for Kisunla.



7. IMPORTANT SAFETY INFORMATION FOR Kisunla™ (donanemab-azbt) (Continued)

Amyloid-Related Imaging Abnormalities (ARIA) (Continued)

In Study 2, symptomatic ARIA-E occurred in 3% of patients and symptomatic ARIA-H occurred in less than 1% of patients through 12 months of treatment with Kisunla. Clinical symptoms associated with ARIA-E resolved in approximately 67% of patients at 12 months. Including asymptomatic radiographic events, ARIA, ARIA-E, and ARIA-H were observed in 29%, 16%, and 25% of patients treated with Kisunla.

Incidence of Intracerebral Hemorrhage (ICH)

ICH >1 cm in diameter was reported in 0.5% of patients treated with Kisunla vs 0.2% on placebo in Study 1 and in 1% of patients treated with Kisunla in Study 2. Fatal events of ICH have been observed in patients taking Kisunla.

Risk Factors for ARIA and ICH

ApoE ε4 Carrier Status

The risk of ARIA, including symptomatic and serious ARIA, is increased in ApoE ε4 homozygotes.

Recommendations for management of ARIA do not differ based on ApoE ε4 carrier status. Testing for ApoE ε4 status should be performed prior to initiation of treatment to inform the risk of developing ARIA. An FDA-authorized test for detection of ApoE ε4 alleles is not currently available. Currently available tests may vary in accuracy and design.

Radiographic Findings of Cerebral Amyloid Angiopathy (CAA)

Neuroimaging findings that may indicate CAA include evidence of prior ICH, cerebral microhemorrhage, and cortical superficial siderosis. CAA has an increased risk for ICH. The presence of an ApoE ε4 allele is also associated with CAA.

The baseline presence of at least 2 microhemorrhages or the presence of at least 1 area of superficial siderosis on magnetic resonance imaging (MRI), which may be suggestive of CAA, were identified as risk factors for ARIA. Patients were excluded from enrollment in Study 1 and Study 2 for findings on neuroimaging of prior ICH >1 cm in diameter, >4 microhemorrhages, >1 area of superficial siderosis, severe white matter disease, and vasogenic edema.

Concomitant Antithrombotic or Thrombolytic Medication

In Study 1, baseline use of antithrombotic medication (aspirin, other antiplatelets, or anticoagulants) was allowed. The majority of exposures to antithrombotic medications were to aspirin. The number of events and the limited exposure to non-aspirin antithrombotic medications limit definitive conclusions about the risk of ARIA or ICH in patients taking antithrombotic medications.

One fatal ICH occurred in a patient taking Kisunla in the setting of focal neurologic symptoms of ARIA and the use of a thrombolytic agent in Study 1, and one fatal intracerebral hemorrhage occurred in the setting of ARIA and the use of a thrombolytic agent in Study 2. Because ARIA-E can cause focal neurologic deficits that can mimic an ischemic stroke, treating clinicians should consider whether such symptoms could be due to ARIA-E, and additional caution should be exercised when considering the administration of antithrombotics or a thrombolytic agent (eg, tissue plasminogen activator) to a patient being treated with Kisunla. Advise patients to carry information that they are being treated with Kisunla.

Please see additional Important Safety Information on [page 16](#), and full [Prescribing Information](#), including Boxed Warning regarding ARIA, and [Medication Guide](#) for Kisunla.



7. IMPORTANT SAFETY INFORMATION FOR Kisunla™ (donanemab-azbt) (Continued)

Amyloid-Related Imaging Abnormalities (ARIA) (Continued)

Caution should be exercised when considering the use of Kisunla in patients with factors that indicate an increased risk for ICH and in particular for patients who need to be on anticoagulant therapy or patients with findings on MRI that are suggestive of CAA.

Monitoring and Dose Management Guidelines

Obtain a recent baseline brain MRI prior to initiating treatment and prior to the 2nd, 3rd, 4th, and 7th infusions. Enhanced clinical vigilance for ARIA is recommended during the first 24 weeks of treatment with Kisunla. If a patient experiences symptoms suggestive of ARIA, clinical evaluation should be performed, including MRI if indicated. If ARIA is observed on MRI, careful clinical evaluation should be performed prior to continuing treatment.

Recommendations for dosing in patients with ARIA-E and ARIA-H depend on clinical symptoms and radiographic severity. Depending on ARIA severity, use clinical judgment in considering whether to continue dosing, interrupt treatment, or permanently discontinue Kisunla. See Prescribing Information for additional dosing considerations.

Hypersensitivity Reactions

Hypersensitivity reactions, including anaphylaxis and angioedema, have occurred in patients who were treated with Kisunla. Promptly discontinue the infusion upon the first observation of any signs or symptoms consistent with a hypersensitivity reaction and initiate appropriate therapy.

Infusion-Related Reactions (IRR)

IRRs were observed with Kisunla with the majority occurring within the first 4 infusions. Most IRRs occurred during the infusion or within 30 minutes after completion of the infusion, however some have occurred hours after an infusion. Signs and symptoms of IRRs include chills, erythema, nausea/vomiting, flushing, difficulty breathing/dyspnea, sweating, elevated blood pressure, headache, chest pain, and low blood pressure.

In the event of an IRR, the infusion rate may be reduced, or the infusion may be discontinued, and appropriate therapy initiated as clinically indicated. Consider pre-treatment with antihistamines, acetaminophen, or corticosteroids prior to subsequent dosing.

Adverse Reactions: The most common adverse reactions reported in $\geq 5\%$ of patients treated with Kisunla and $\geq 2\%$ higher than placebo were ARIA-H microhemorrhage, ARIA-E, ARIA-H superficial siderosis, headache, and IRRs.

Kisunla (donanemab-azbt) injection for intravenous use is available as a 350 mg/20 mL single-dose vial.

Please see full [Prescribing Information](#), including Boxed Warning regarding ARIA, and [Medication Guide](#) for Kisunla.

DN HCP ISI 01JUL2025

This image shows a single sheet of white paper with horizontal ruling lines. The lines are evenly spaced and run across the width of the page. There are no margins, text, or other markings on the paper.

 **kisunla**[™]
(donanemab-azbt) | injection for
intravenous use
350mg/20mL

A Lilly Medicine

For more information about Kisunla, visit <https://kisunla.lilly.com/hcp>

Please see additional Important Safety Information on **pages 14-16**, and full **Prescribing Information**, including Boxed Warning regarding ARIA, and **Medication Guide** for Kisunla.

REFERENCE: 1. Kisunla (donanemab-azbt) Prescribing Information. Lilly USA, LLC.

Oracle Health is a trademark of Oracle.

Kisunla™ is a trademark owned or licensed by Eli Lilly and Company, its subsidiaries, or affiliates.

PP-DN-US-0564 07/2025 © Lilly USA, LLC 2025. All rights reserved.



A Lilly Medicine

